arXiv:1001.1905v2 [math.CO] 30 May 2011

BOUNDS ON SOME EDGE FOLKMAN NUMBERS [*]

May 31, 2011

Nikolay R. Kolev

Abstract

The edge Folkman numbers are defined by the equality

$$F_e(a_1, \dots, a_r; q) = \min\{|V(G)| : G \overset{e}{\rightarrow} (a_1, \dots, a_r; q) \text{ and } cl(G) < q\}.$$

Lin proved that if $R(a_1, a_2) = R(a_1-1, a_2)+R(a_1, a_2-1)$ then $F_e(a_1, a_2; R(a_1, a_2)) = R(a_1, a_2)+2$, where $R(a_1, a_2)$ is the Ramsey number showing that $K_{R(a_1,a_2)-3}+C_5 \overset{e}{\rightarrow} (a_1, a_2)$ where C_5 is the simple cycle on 5 vertices. We prove some upper bounds on edge Folkman numbers for which $R(a_1, a_2) < R(a_1 - 1, a_2) + R(a_1, a_2 - 1)$ and we cite some lower bounds.

1 1. Introduction

Only finite non-oriented graphs without multiple edges and loops are considered. We call a p-clique of the graph G a set of p vertices each two of which are adjacent. The largest positive integer p such that G contains a p-clique is denoted by cl(G). A set of vertices of the graph G none two of which are adjacent is called an independent set. The largest positive integer p such that G contains an independent set on p vertices is called the independence number of the graph G and is denoted by $\alpha(G)$. In this paper we shall also use the following notations:

- $V(G)$ is the vertex set of the graph G;
- $E(G)$ is the edge set of the graph G;
- $N(v)$, $v \in V(G)$ is the set of all vertices of G adjacent to v;
- $G[V]$, $V \subseteq V(G)$ is the subgraph of G induced by V;
- K_n is the complete graph on n vertices;
- $\overline{G}$ is the complementary graph of G.

---

[*] Partially supported by the scientific research fund of St. Kl. Ohridski Sofia University under contract No 75/2009

Let G_1 and G_2 be two graphs without common vertices. We denote by $G_1 + G_2$ the graph G for which $V(G) = V(G_1) \cup V(G_2)$ and $E(G) = E(G_1) \cup E(G_2) \cup E'$ where $E' = \{xy : x \in V(G_1), y \in V(G_2)\}$. It is clear that

$$cl(G_1 + G_2) = cl(G_1) + cl(G_2). \tag{1}$$

Definition 1. Let $a_1, \ldots, a_r$ be positive integers. The symbol $G \overset{v}{\to} (a_1, \ldots, a_r)$ means that in every r-coloring of $V(G)$ there is a monochromatic a_i-clique in the i-th color for some $i \in \{1, \ldots, r\}$.

Definition 2. Let $a_1, \ldots, a_r$ be positive integers. We say that an r-coloring of $E(G)$ is $(a_1, \ldots, a_r)$-free if for each $i = 1, \ldots, r$ there is no monochromatic a_i-clique in the i-th color. The symbol $G \overset{e}{\to} (a_1, \ldots, a_r)$ means that there is no $(a_1, \ldots, a_r)$-free coloring of $E(G)$.

The smallest positive integer n for which $K_n \overset{e}{\to} (a_1, \ldots, a_r)$ is called a Ramsey number and is denoted by $R(a_1, \ldots, a_r)$. Note that the Ramsey number $R(a_1, a_2)$ can be interpreted as the smallest positive integer n such that for every n-vertex graph G either $cl(G) \geq a_1$ or $\alpha(G) \geq a_2$. The existence of such numbers was proved by Ramsey in [18]. We shall use only the values $R(3, 3) = 6$ and $R(3, 4) = 9$, [3].

The edge Folkman numbers are defined by the equality

$$F_e(a_1, \ldots, a_r; q) = \min\{|V(G)| : G \overset{e}{\to} (a_1, \ldots, a_r) \text{ and } cl(G) < q\}.$$

It is clear that $G \overset{e}{\to} (a_1, \ldots, a_r)$ implies $cl(G) \geq \max\{a_1, \ldots, a_r\}$. There exists a graph G such that $G \overset{e}{\to} (a_1, \ldots, a_r)$ and $cl(G) = \max\{a_1, \ldots, a_r\}$. In the case $r = 2$ this was proved in [1] and in the general case in [16]. Therefore

$$F_e(a_1, \ldots, a_r; q) \text{ exists if and only if } q > \max\{a_1, \ldots, a_r\}.$$

It follows from the definition of $R(a_1, \ldots, a_r)$ that

$$F_e(a_1, \ldots, a_r; q) = R(a_1, \ldots, a_r) \text{ if } q > R(a_1, \ldots, a_r).$$

The smaller the value of q in comparison to $R(a_1, \ldots, a_r)$ the more difficult the problem of computing the number $F_e(a_1, \ldots, a_r; q)$. Only ten edge Folkman numbers that are not Ramsey numbers are known. For the results see the papers: [2],[4], [6], [7], [9], [10], [11], [13].

2 2. Upper bounds on the edge Folkman numbers

We obtained the following result:

Theorem 1 *Let* a *and* α *be nonnegative integers. Let us denote* $R = R(3, a)$. *Let* $R = R(3, a-1) + a - \alpha$ *and* $R - 3a + \alpha + 5 \geq R(3, a-2)$, $a \geq 4$. *If there exists a graph* U *with the properties*

$$cl(U) = a - 1$$

$$U \overset{v}{\to} (a-1, a-2)$$

$$U \overset{v}{\to} (\underbrace{a-3, \ldots, a-3}_{a-2 \text{ times}}, 3).$$

Then

$$F_e(3,a;R-a+\alpha+4) \le R-2a+\alpha+4+|V(U)|.$$

Proof Consider the graph $G = K_{R-2a+\alpha+4} + U$ where U is the graph from the statement of the theorem. We will prove that $G \overset{e}{\to} (3,a)$ and thus the theorem will be proved (here we use (1) to compute that $cl(G) = R-2a+\alpha+4+cl(U) = R-a+\alpha+3$).

Assume the opposite: that there exists a $(3,a)$-free 2-coloring of $E(G)$

$$E(G) = E_1 \cup E_2, \qquad E_1 \cap E_2 = \emptyset. \tag{2}$$

We shall call the edges in E_1 blue and the edges in E_2 red.

We define for an arbitrary vertex $v \in V(G)$ and index $i = 1, 2$:

$$\begin{aligned} &N_i(v) = \{x \in N(v) \mid [v,x] \in E_i\},\\ &G_i(v) = G[N_i(v)]\\ &A_i(v) = N_i(v) \cap V(U) \end{aligned}$$

Let H be a subgraph of G. We say that H is a monochromatic subgraph in the blue-red coloring (2) if $E(H) \subseteq E_1$ or $E(H) \subseteq E_2$. If $E(H) \subseteq E_1$ we say that H is a blue subgraph and if $E(H) \subseteq E_2$ we say that H is a red subgraph.

It follows from the assumption that the coloring (2) is $(3,a)$-free that

$$cl(G_1(v)) \le a-1 \text{ and } cl(G_2(v)) \le R(3,a-1)-1 \text{ for each } v \in V(G) \tag{3}$$

Indeed, assume that $cl(G_1(v)) \ge a$. Then there must be no blue edge connecting any two of the vertices in $G_1(v)$ because otherwise this blue edge together with the vertex v would give a blue triangle. As we assumed $cl(G_1(v)) \ge a$ then we have a red a-clique. Analogously assume $cl(G_2(v)) \ge R(3,a-1)$. Then we have either a blue 3-clique or a red $(a-1)$-clique in $G_2(v)$. If we have a blue 3-clique in $G_2(v)$ then we are through. If we have a red $(a-1)$-clique then this $(a-1)$-clique together with the vertex v gives a red a-clique. Thus (3) is proved.

We shall prove that

$$cl(G[A_1(v)]) + cl(G[A_2(v)]) \le 2a-5 \text{ for each } v \in V(K_{R-2a+\alpha+4}) \tag{4}$$

Assume that (4) is not true, i.e. that there exists a vertex $v \in V(K_{R-2a+\alpha+4})$ such that

$$cl(G[A_1(v)]) + cl(G[A_2(v)]) \ge 2a-4.$$

Then as there are $R-2a+\alpha+3$ more vertices in $V(K_{R-2a+\alpha+4})$ with the exception of v, it follows that

$$cl(G_1(v)) + cl(G_2(v)) \ge R-2a+\alpha+3+2a-4 = R+\alpha-1 = R(3,a-1)+a-1$$

(here we use the statement of the theorem that $R(3,a) = R(3,a-1)+a-\alpha$.)

This contradicts (3). Thus (4) is proved.

Now we shall prove that

$$cl(G[A_1(v)]) = a-1 \text{ or } cl(G[A_2(v)]) = a-1 \text{ for each } v \in V(K_{R-2a+\alpha+4}) \tag{5}$$

Assume that (5) is not true. Then we obtain from $cl(U) = a-1$ that

$$cl(G[A_1(v)]) \le a-2 \text{ and } cl(G[A_2(v)]) \le a-2 \text{ for some } v \in V(K_{R-2a+\alpha+4}). \tag{6}$$

It follows from the statement of the theorem that $U \overset{v}{\rightarrow} (a-1, a-2)$ that in every 2-coloring of $V(U)$, in which there are no $(a-1)$-cliques in none of the two colors then there are $(a-2)$-cliques in the both colors.Hence $A_1(v)$ and $A_2(v)$ contain $(a-2)$-clique. Therefore the inequalities in (6) are in fact equalities, which contradicts (4). Thus (5) is proved. Let us note that it follows from (5) and (4) that for each $v \in V(K_{R-2a+\alpha+4})$

$$\text{if } cl(G[A_2(v)]) = a-1 \text{ then } cl(G[A_1(v)]) \leq a-4. \tag{7}$$

If we assume that there are a vertices v in $V(K_{R-2a+\alpha+4})$ with the property $cl(G[A_1(v)]) = a-1$ then it follows from (3) that there are only red edges between these a vertices and hence this is a red a-clique. Therefore having in mind the statement of the theorem there are at least $R-2a+\alpha+4-(a-1) \geq R(3, a-2)$ vertices v in $V(K_{R-2a+\alpha+4})$ with the property $cl(G[A_1(v)]) \leq a-2$. Let us denote the set of these vertices by S. It follows by (5) that $cl(G[A_2(v)]) = a-1$ for each $v \in S$. Then we obtain from (7) that

$$cl(G[A_1(v)]) \leq a-4 \text{ for each } v \in S. \tag{8}$$

As there are no blue 3-cliques among the vertices in S there is a red $(a-2)$-clique among them. Let us denote the vertices of this $(a-2)$-clique by w_i, $i \in 1, \ldots, a-2$. Let us partition the vertices of U in $a-1$ colors in the following way:

$$V_1 = A_1(w_1)$$

and for $j \in 2, \ldots, a-2$

$$V_j = A_1(w_j) \backslash V_1 \cup \ldots \cup V_{j-1}$$

$$V_{a-1} = V(U) \backslash V_1 \cup \ldots \cup V_{a-2}.$$

According to (8) it is impossible V_j, $j \in 1, \ldots, a-2$ to contain a $(a-3)$-clique because according to its definition V_j is a subset of $A_1(w_j)$. But we know from the statement of the theorem that $U \overset{v}{\rightarrow} (\underbrace{a-3, \ldots, a-3}_{a-2 \text{ times}}, 3)$. Hence V_{a-1} contains a 3-clique. Then this 3-clique must contain a red edge (otherwise it is a blue 3-clique and we are through). But it follows from the definition of V_{a-1} that all the edges between the vertices in V_{a-1} and the vertices $w_1, \ldots, w_{a-2}$ are red. Then the red edge in V_{a-1} and the red $(a-2)$-clique $w_1, \ldots, w_{a-2}$ form a red $a-$clique. Thus the theorem is proved.

If we put $\alpha = 0$, $a = 5$, $U = Q$, where Q denotes the graph whose complementary is the graph given here we obtain the following Kolev‘s result

$$F_e(3, 5; 13) \leq 21, [5].$$

What is novel in this result is that for the first time the graph Q is used in the theory of edge Folkman numbers.

We also prove the following result which unfortunately is not a particular case of **Theorem1**:

Theorem 2 $F_e(4, 4; 17) \leq 25$.

Proof Consider the graph $G = K_{12} + Q$ and we shall show that $G \overset{e}{\rightarrow} (4, 4)$.

Assume the opposite: that there exists a $(4, 4)$-free 2-coloring of $E(G)$

$$E(G) = E_1 \cup E_2, \qquad E_1 \cap E_2 = \emptyset. \tag{9}$$



Figure 1: Graph $\overline{Q}$

We shall call the edges in E_1 blue and the edges in E_2 red.

We define for an arbitrary vertex $v \in V(G)$ and index $i = 1, 2$:

$$\begin{aligned} N_i(v) &= \{x \in N(v) \mid [v, x] \in E_i\}, \\ G_i(v) &= G[N_i(v)] \\ A_i(v) &= N_i(v) \cap V(Q) \end{aligned}$$

We say that H is a monochromatic subgraph in the blue-red coloring (9) if $E(H) \subseteq E_1$ or $E(H) \subseteq E_2$. If $E(H) \subseteq E_1$ we say that H is a blue subgraph and if $E(H) \subseteq E_2$ we say that H is a red subgraph.

It follows from the assumption that the coloring (9) is (4,4)-free and $R(3, 4) = 9$ that

$$cl(G_1(v)) \leq 8 \text{ and } cl(G_2(v)) \leq 8 \text{ for each } v \in V(G) \tag{10}$$

Assume $cl(G_2(v)) \geq 9$. Since $R(3, 4) = 9$, then we have either a blue 3-clique or a red 4-clique in $G_1(v)$. If we have a blue 3-clique in $G_1(v)$ then this blue 3-clique together with the vertex v forms a blue 4-clique. If we have a red 4-clique then we are through. Analogously we disprove $cl(G_1(v)) \geq 9$. Thus (10) is proved.

We shall prove that

$$cl(G[A_1(v)]) + cl(G[A_2(v)]) \leq 5 \text{ for each } v \in V(K_{12}) \tag{11}$$

Assume that (11) is not true, i.e. that there exists a vertex $v \in V(K_{12})$ such that

$$cl(G[A_1(v)]) + cl(G[A_2(v)]) \geq 6.$$

Then as there are eleven more vertices in $V(K_{12})$ with the exception of v, it follows that

$$cl(G_1(v)) + cl(G_2(v)) \geq 17.$$

It follows from the pigeonhole principle that either $cl(G_1(v)) \geq 9$ or $cl(G_2(v)) \geq 9$, which contradicts (10). Thus (11) is proved.

It follows from $Q \overset{e}{\to} (3,4)$ and $cl(Q) = 4$ that

$$cl(G[A_1(v)]) = 4 \text{ or } cl(G[A_2(v)]) = 4 \text{ for each } v \in V(K_{12}). \tag{12}$$

It follows from (11) and (12) that

$$\text{if } cl(G[A_1(v)]) = 4 \text{ then } cl(G[A_2(v)]) \leq 1 \text{ for each } v \in V(K_{12}). \tag{13}$$

Assume that there are two vertices $a, b \in V(K_{12})$ such that the edge ab is blue and $cl(G[A_1(a)]) = cl(G[A_1(b)]) = 4$. Then it follows from (13) that $cl(G[A_2(a)]) \leq 1$ and $cl(G[A_2(b)]) \leq 1$. Let us consider the following 3-coloring of $V(Q)$:

$$V_1 = A_2(a)$$

$$V_2 = A_2(b) \backslash A_1(a)$$

$$V_3 = V(Q) \backslash (V_1 \cup V_2).$$

It is clear from the definition of V_1, V_2, V_3 that V_3 consists of such vertices that are connected with both a and b with blue edges. It is clear from $cl(G[A_2(a)]) \leq 1$ and $cl(G[A_2(b)]) \leq 1$ that V_1 and V_2 contain no 2-cliques. But we know from [8] that $Q \overset{v}{\to} (2,2,4)$ and hence V_3 contains a 4-clique. If this 4-clique has a blue edge then then this blue edge together with the edge ab is a blue 4-clique. If this 4-clique does not have a blue edge then it is a red 4-clique- a contradiction. Thus we obtained that

$$\text{if } cl(G[A_1(a)]) = cl(G[A_1(b)]) = 4 \text{ then the edge } ab \text{ is not blue, } a, b \in K_{12}. \tag{14}$$

Analogously we obtain

$$\text{if } cl(G[A_2(a)]) = cl(G[A_2(b)]) = 4 \text{ then the edge } ab \text{ is not red, } a, b \in K_{12}. \tag{15}$$

We have from (12) that $cl(G[A_1(v)]) = 4$ or $cl(G[A_2(v)]) = 4$ for each $v \in K_{12}$. For a fixed $v \in K_{12}$ we may assume without loss of generality that $cl(G[A_1(v)]) = 4$. Assume that for this v there exist three distinct vertices $b_1, b_2, b_3 \in K_{12}$ such that the edges vb_i, $i \in \{1,2,3\}$ are blue. Now it follows from (13), (14),(15) and the fact that the edges vb_i, $i \in \{1,2,3\}$ are blue that

$$cl(G[A_2(b_i)]) = 4 \text{ and } cl(G[A_1(b_i)]) \leq 1 \tag{16}$$

for $i \in \{1,2,3\}$. Now it follows from (15) that the edges b_ib_j are blue for $i, j \in \{1,2,3\}$. Hence $vb_1b_2b_3$ is a blue 4−clique and the theorem is proved.

Thus we obtained that there are no more than two blue edges from the vertex v. Therefore there are at least 9 vertices connected to v with red edges which contradicts (10). We proved that $G \overset{e}{\to} (4,4)$. As $cl(G) = 4$ then we obtain that $F_e(4,4;17) \leq 25$, which we wanted to prove.

Remark So far it was known by [1] that $F_e(4,4;17) < \infty$.

The last known lower bound of the edge Foklman number $F_e(4,4;17)$ is $F_e(4,4;17) \geq 22$ - see [15].

3 References

……

[1]Folkman, J. Graphs with monochromatic complete subgraphs in every edge coloring. *SIAM J. Appl. Math.* **18**, 1970, 19–24. [2] Graham R.L. On edgewise 2-colored graphs with monochromatic triangles containing no complete hexagon *J. Comb. theory* **4**, 1968, 300. [3]Greenwood, R., A. Gleason. Combinatorial relation and chromatic graphs. *Canad. J. Math., **7**, 1955, 1–7.* [4] Lin S. On Ramsey numbers and K_r-coloring of graphs *J. Comb. theory Ser B* **12**, 1972, 82-92. [5] Kolev N. New upper bound for the edge Folkman number $F_e(3,5;13)$. *Serdica math. J., **34** 2008,783-790.* [6]Kolev, N., N. Nenov. An example of 16-vertex Folkman edge (3,4)-graph without 8-cliques. *Annuaire Univ. Sofia Fac. Math. Inform., **98**, 2008 101- 126, see http* $: //arxiv.org/PS_cache/math/pdf/0602/0602249v1.pdf$. [7] Kolev N., N. Nenov. The Folkman number $F_e(3,4;8)$ is equal to 16. *Compt. rend. Acad. bulg. Sci., **59**,No 1, 2006, 25–30.* [8]Nenov N.On the vertex Folkman number F(3,4). *Compt. rend. Acad. bulg. Sci., **54**, 2, 2001, 23-26.* [9] Nenov N. On an assumption of Lin about Ramsey-Graham-Spencer numbers. *(Russian)Compt. rend. Acad. bulg. Sci., **33**, 9, 1980, 1171-1174.* [10] Nenov N. Generalization of a certain theorem of Greenwood and Gleason on three-color colorings of the edges of a complete graph with 17 vertices. *(Russian)Compt. rend. Acad. bulg. Sci., **34**, 1981, 1209-1212.* [11] Nenov N. Lower estimates for some constants related to Ramsey graphs.*(Russian) Annuaire Univ. Sofia Fac. Math. Inform., **75**, 1984, 27-38.* [12]Nenov N. On the (3,4)-Ramsey graphs without 9-cliques. *(Russian) Univ. Sofia Fac. Math. Inform., **85**, 1991 ,71-81 .* [13] Nenov N. An example of 15-vertex Ramsey (3,3)-graph with clique number 4. *(Russian)Compt. rend. Acad. bulg. Sci., **34**, 1981, 1487-1489.* [14] Nenov N.On the Zykov numbers and some of its applications to Ramsey theory. *(Russian)Compt. rend. Acad. bulgSerdica Bulg. math. Publ.9,161-167, (1983).* [15] Nenov N. On the vertex Folkman numbers$F_v(2,\dots,2;q)$. *Serdica Math. J. 35, 2009 251-272.* [16]Nesetril J, Rodl V. The Ramsey property for graphs with forbidden complete subgraphs.*J. Comb Th. **B20**, 1976, 243-249.* [17] Piwakowski K., Radziszowski S., Urbanski S. Computation of the Folkman number $F_e(3,3;5)$. *J. Graph theory* **32**, 1999, 41-49. [18]Ramsey P. On a problem of formal logic *Proc. London Math. Soc., 30, 1930, 264-286.*

Faculty of mathematics and informatics
Sofia University
Sofia 1164, BULGARIA
E-MAIL: nickyxy@fmi.uni-sofia.bg

Received xx.xx.199x

This figure "01.jpg" is available in "jpg" format from:

http://arxiv.org/ps/1001.1905v2